



New Cases 

Age-adjusted rate per 100,000 people

Top 5		
1	Utah	26.6
2	New Mexico	36.2
3	California	39.7
4	Colorado	39.7
5	Wyoming	42.0
Bottom 5		
47	Mississippi	73.2
48	Tennessee	73.3
49	Arkansas	75.8
50	West Virginia	78.3
51	Kentucky	87.6

Survival 

Percent of cases still alive five years after diagnosis

Top 5		
1	Rhode Island	30.8%
2	Connecticut	30.5%
3	New York	29.9%
4	Minnesota	29.7%
5	Oregon	28.5%
5	New Jersey	28.5%
Bottom 5		
42	Mississippi	20.4%
42	West Virginia	20.4%
42	Kentucky	20.4%
45	Alabama	20.0%
46	Oklahoma	19.7%

Early Diagnosis 


Percent of cases diagnosed at an early stage

Top 5		
1	Massachusetts	31.9%
2	Connecticut	31.3%
3	Rhode Island	30.4%
4	New York	29.9%
5	Wyoming	29.1%
Bottom 5		
45	Louisiana	22.9%
46	Alaska	22.5%
47	Oklahoma	22.1%
48	New Mexico	20.7%
49	Hawaii	19.5%

Surgery 


Percent of cases undergoing surgery as part of their first course of treatment

Top 5		
1	Massachusetts	31.2%
2	New York	29.1%
3	Rhode Island	28.3%
4	New Jersey	26.1%
5	Connecticut	25.7%
Bottom 5		
45	Texas	16.1%
46	South Dakota	16.0%
47	Oklahoma	15.7%
48	Wyoming	15.3%
49	New Mexico	13.5%

Untreated 

Percent of cases receiving no treatment after being diagnosed

Top 5		
1	North Dakota	13.9%
2	Massachusetts	14.0%
2	Rhode Island	14.0%
4	Missouri	15.4%
5	Ohio	16.4%
Bottom 5		
44	Florida	25.1%
44	Hawaii	25.1%
46	California	25.8%
47	Texas	28.4%
48	New Mexico	29.7%
49	Arizona	31.7%

Screening 


Percent of those at high risk receiving screening

Top 5		
1	Massachusetts	16.3%
2	Rhode Island	13.4%
3	Vermont	13.2%
4	Kentucky	13.0%
5	Maine	11.8%
Bottom 5		
45	Utah	2.0%
45	Oklahoma	2.0%
45	New Mexico	2.0%
48	Texas	1.9%
49	Wyoming	1.7%
50	Nevada	1.3%
51	California	1.0%

Smoking 

Percent of adults who are current smokers

Top 5		
1	Utah	8.2%
2	California	8.9%
3	New Jersey	10.8%
4	Maryland	10.9%
5	Massachusetts	11.1%
Bottom 5		
47	Tennessee	19.5%
48	Mississippi	20.1%
49	Arkansas	20.5%
50	Kentucky	21.4%
51	West Virginia	22.6%

Radon 

Estimated Percent of Radon Test Results ≥4 pCi/L (EPA Action Level)

Top 5		
1	Hawaii	0.4%
2	Louisiana	3.1%
3	Mississippi	5.2%
4	Texas	7.7%
5	California	8.2%
Bottom 5		
47	Ohio	48.8%
48	Nebraska	55.8%
49	Iowa	57.9%
50	North Dakota	58.0%
51	South Dakota	59.8%